

Article

Lower extremity kinetics and kinematics in runners with patellofemoral pain: a retrospective case-control study using musculoskeletal simulation

Sinclair, Jonathan Kenneth, Chockalingam, Nachi and Taylor, Paul John

Available at <http://clock.uclan.ac.uk/40321/>

Sinclair, Jonathan Kenneth ORCID: 0000-0002-2231-3732, Chockalingam, Nachi and Taylor, Paul John ORCID: 0000-0002-9999-8397 (2022) Lower extremity kinetics and kinematics in runners with patellofemoral pain: a retrospective case-control study using musculoskeletal simulation. Applied Sciences, 12 (2).

It is advisable to refer to the publisher's version if you intend to cite from the work.
<http://dx.doi.org/10.3390/app12020585>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.

Lower extremity kinetics and kinematics in runners with patellofemoral pain: a retrospective case-control study using musculoskeletal simulation.

Jonathan Sinclair^{1*}, Nachiappan Chockalingam² & Paul John Taylor^{1,3}

¹ Research Centre for Applied Sport, Physical Activity and Performance, School of Sport & Health Sciences, Faculty of Allied Health and Wellbeing, University of Central Lancashire, Lancashire, UK.

² School of Life Sciences and Education, Staffordshire University, City of Stoke, Stoke on Trent, UK.

³ School of Psychology and Computer Sciences, Faculty of Science and Technology, University of Central Lancashire, Lancashire, UK.

* Correspondence: jksinclair@uclan.ac.uk

Abstract: Patellofemoral pain (PFP) is a common atraumatic knee pathology in runners, with a complex multifactorial aetiology influenced by sex differences. This retrospective case-control study therefore aimed to evaluate lower limb kinetics and kinematics in symptomatic and control male and female runners using musculoskeletal simulation. Lower extremity biomechanics were assessed in 40 runners with PFP (15 females & 25 males) and 40 controls (15 females & 25 males), whilst running at a self-selected velocity. Lower extremity biomechanics were explored using a musculoskeletal simulation approach. Four between group comparisons 1. Overall PFP vs. Control, 2. Male PFP vs. Male Control, 3. Female PFP vs. Female Control and 4. Male PFP vs. Female PFP were undertaken using linear mixed models. The overall (stress per mile: PFP=1047.49 & Control=812.93) and female (peak stress: PFP=13.07KPa/BW & Control=10.82KPa/BW) comparisons showed increased patellofemoral joint stress indices in PFP runners. A significantly lower strike index was also shown in PFP runners in the overall (PFP=17.75% & Control=33.57%) and female analyses (PFP=15.49% & Control=40.20%), which revealed a midfoot strike in control, and a rearfoot pattern in PFP runners. Peak rearfoot eversion and contralateral pelvic drop range of motion (ROM) were shown to be greater in PFP runners in the overall (eversion: PFP=−8.15° & Control=−15.09°/ pelvic drop ROM: PFP=3.64° & Control=1.88°), male (eversion: PFP=−8.05° & Control=−14.69°/ pelvic drop ROM: PFP=3.16° & Control=1.77°) and female (eversion: PFP=8.28° & Control=−15.75°/ pelvic drop ROM: PFP=3.64° & Control=1.88°) PFP runners, whilst female PFP runners (11.30°) exhibited a significantly larger peak hip adduction compared to PFP males (7.62°). The findings from this investigation identify biomechanical differences between control and PFP runners as well as demonstrating distinctions in PFP presentation for many parameters between sexes. Highlighting potential risk factors for PFP that may be addressed through focused intervention modalities and also the need where appropriate, for sex specific targeted treatment approaches.

Keywords: Patellofemoral pain; Running; Musculoskeletal simulation; Anterior knee pain; Kinematics.

1. Introduction

Both recreational and competitive distance running is associated with a plethora of physiological benefits [1]. However, despite the physical improvements mediated by running, it is also linked with a very high rate of overuse injuries [2, 3]. Patellofemoral pain (PFP) is a common atraumatic knee pathology that typically manifests as retropatellar or peripatellar pain and inflammation, aggravated by activities that frequently load the joint [4]. PFP has an incidence rate as high as 21% in runners [5], with a much larger prevalence in females [6]. The long-term prognosis is poor, with between 71–91% of patients experiencing symptoms 20 years following diagnosis [7]. PFP symptoms may later present with radiographic evidence of osteoarthritis at this joint [8], and pain symptoms force many to

Citation: Lastname, F.; Lastname, F.; Lastname, F. Title. *Appl. Sci.* **2021**, *11*, x. <https://doi.org/10.3390/xxxxx>
Academic Editor: Firstname Lastname

Received: date

Accepted: date

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

reduce or even end their running training [9]; causing many to develop associated psychological disorders [10].

The lack of treatment efficacy is attributed to the multifactorial nature and complex aetiology of PFP, which makes the identification of modifiable risk factors problematic [11]. Given the high incidence of PFP in runners [5]; an enhanced comprehension of the parameters linked to PFP aetiology and its distinct incidence in differing populations is essential to prevent symptoms and improve management. Recent prospective pooled systematic review and meta-analytic analyses have shown physiological indices of quadriceps weakness and hip abduction strength to be risk factors for PFP in military and adolescent populations, although no identifiable risk factors were evident in runners [12].

There are a lack of prospective investigations exploring the biomechanical mechanisms linked to the aetiology of PFP in runners [4]. Noehren et al., [13], showed that female runners were at an increased risk of developing PFP in the presence of a statistically greater maximum hip adduction angle. Furthermore, in relation to asymptomatic controls, Fox et al., [14] showed that PFP runners exhibited increased peak hip adduction, hip transverse plane range of motion (ROM), peak knee flexion, peak knee abduction and peak dorsiflexion. Noehren et al., [15, 16] revealed that in female PFP runners, increased hip adduction, hip internal rotation and tibial internal rotation was evident compared to controls. Esculier et al., [17] showed that runners with PFP exhibited higher hip adduction at toe-off, lower hip adduction ROM during late-stance and longer of soleus activation. Conversely Dierks et al., [18] showed that runners with PFP demonstrated reduced peak knee flexion, peak hip adduction, eversion ROM, peak knee flexion velocity, peak hip adduction velocity, and peak hip internal rotation velocity compared with controls. Finally, Willson & Davis, [19] revealed that females with PFP demonstrated greater hip adduction and contralateral pelvic drop in relation to asymptomatic runners. Neal et al., [12] proposed that these kinematic variations contribute to the development and persistence of PFP by way of increased patellofemoral joint stress, although patellofemoral joint loading has not yet been explored in runners with PFP in relation to asymptomatic controls.

Although previous analyses have explored running biomechanics in those with PFP compared to healthy controls, the majority utilized only female runners or a mixed sex sample, without considering of the effects of sex. As such the effects of sex on running biomechanics in those with PFP is poorly understood [12]. Willy et al., [20] compared running biomechanics in male and female runners with PFP and showed that females exhibited greater peak hip adduction compared to both males with PFP and male controls, whereas males with PFP ran with a greater peak knee adduction compared to symptomatic females and male controls. A drawback of this investigation was the lack of a female control group and Neal et al., [4] undertook the only investigation examine male and female runners with PFP in relation to asymptomatic controls of both sexes. Mixed-sex comparisons showed that PFP runners exhibited greater peak hip adduction compared to controls. Females with PFP demonstrated significantly greater peak hip adduction compared to controls but there were no differences between males with PFP and male controls.

Previous analyses concerning the biomechanical differences between runners with PFP and asymptomatic controls have examined only joint kinematics, ground reaction forces and surface electromyography. Significant advances have been made in musculoskeletal simulation models [21], leading to the development of bespoke software, allowing skeletal muscle forces to be simulated during movement and utilized as inputs to calculate lower extremity joint reaction forces [22]. Such approaches have been used extensively in running, but only one study has explored differences in those with PFP and healthy controls. Besier et al., [23] showed that medial gastrocnemius forces were significantly larger in PFP runners, which they speculated may cause greater joint contact loads in comparison to pain-free runners, although joint loading indices were not undertaken in this investigation. With the high incidence of PFP in runners [5], allied to a poor long-term prognosis [7], further investigation is warranted to better understand the biomechanical variables

associated with PFP and the increased incidence of this pathology in female runners, to enhance the quality of future evidence based preventative strategies for both sexes.

This investigation aimed via a retrospective case-control study design, to firstly evaluate running biomechanics in a pooled sex sample of runners with and without PFP using a musculoskeletal simulation-based approach. Secondly using the same experimental approach, this investigation seeks to explore differences between PFP and asymptomatic controls, when these cohorts were divided into subset comparisons of males and females with and without PFP, to investigate distinctions in PFP presentation between the sexes.

2. Materials and Methods

2.1 Participants

The University of Central Lancashire granted ethical approval for this study (STEMH 424), and all participants provided written informed consent prior to participation in accordance with the Declaration of Helsinki. Runners with and without PFP were recruited from local sports physiotherapy clinics and running clubs/ teams. An a priori sample size calculation for independent group comparisons between sexes was undertaken, using the formulae outlined by Rosner, [24]. The peak patellofemoral joint stress was adopted as the primary dependent variable. Using data from our previous work, (males with PFP 13.10 ± 3.05 megapascals (MPa) [25], male controls $10.80 \text{ MPa} \pm 3.04$ [26], females with PFP $10.60 \text{ MPa} \pm 1.20$ [27], and female controls $9.27 \text{ MPa} \pm 1.36$ [28]) it was determined that to achieve $\alpha = 5\%$ and $\beta = 0.80$, 15 female and 25 male runners were required in both the control and PFP groups.

Specific diagnosis of PFP was made in accordance with previous guidelines [29], it was also required that runners had experienced patellofemoral symptoms for a minimum of 3 months that were exacerbated by running or other activities. Participants with any other knee/ lower limb pathology or previous lower limb surgery were excluded. For inclusion in the control group, it was necessary for runners to be free of running-related injury for at least three months and have no prior history of anterior knee pain. Both groups were commonly; participants of either sex, running at least of 10 km per week and aged between 18–45 years [4].

2.2 Procedures

Participants presented to the Biomechanics Laboratory at the University of Central Lancashire. In runners who experienced bilateral PFP, the limb that exhibited the greater symptoms was included [4]. For the control participants, the dominant limb (defined as the limb that would be used to kick a football) was utilized [20]. Participants in the PFP group completed the Knee Osteoarthritis Outcome Score Patellofemoral subscale (KOOS-PF) This scale is scored from 0 to 100, with 100 representing no disability and 0 representing maximum disability [30].

Participants ran across the biomechanics laboratory, striking an embedded piezoelectric force platform (Kistler Instruments Ltd., Winterthur, Switzerland) which sampled at 1000 Hz, with their appropriate foot. Participants were required to run in their usual running shoes and at their own typical/ self-selected running speed. The stance phase was delineated as the duration over which 20 N or greater of vertical ground reaction force (GRF) was applied to the force platform [31]. Runners completed five successful trials, with a successful trial being defined as one where the foot made full contact with the force platform and with no evidence of gait modifications due to the experimental conditions. Kinematic and GRF data were synchronously collected. Kinematic data were captured at 250 Hz via an eight-camera motion analysis system (Qualisys Medical AB, Goteburg, Sweden). Dynamic calibration of the motion capture system was performed before each data collection session.

Body segments were modelled in 6 degrees of freedom using the calibrated anatomical systems technique [32]. To define the anatomical frames of the thorax, pelvis, thighs, shanks and feet, retroreflective markers were placed at the seventh cervical vertebrae (C7),

twelfth thoracic vertebrae (T12) and xiphoid process landmarks, and also positioned bilaterally onto the acromion process, iliac crest, anterior superior iliac spine (ASIS), posterior superior iliac spine (PSIS), medial and lateral malleoli, medial and lateral femoral epicondyles, greater trochanter, calcaneus, first metatarsal and fifth metatarsal (Figure 1a). Intra-rater reliability for the researcher responsible for positioning of the aforementioned anatomical markers has been shown to be excellent [33]. The centre's of the ankle and knee joints were delineated as the mid-point between the malleoli and femoral epicondyle markers [34, 35]. The hip joint centre was determined using a regression equation that uses the positions of the ASIS markers [36]. Carbon-fibre tracking clusters comprising of four non-linear retroreflective markers were positioned onto the thigh and shank segments. In addition, the foot segments were tracked via the calcaneus, first metatarsal and fifth metatarsal, the pelvic segment was tracked using the PSIS and ASIS markers and the thorax segment was tracked using the T12, C7 and xiphoid markers. Static calibration trials were obtained allowing for the anatomical markers to be referenced in relation to the tracking markers/clusters. The Z (transverse) axis was oriented vertically from the distal segment end to the proximal segment end. The Y (coronal) axis was oriented in the segment from posterior to anterior. Finally, the X (sagittal) axis orientation was determined using the right-hand rule and was oriented from medial to lateral (Figure 1b).

@@@FIGURE 1 NEAR HERE@@@

2.3 Processing

Dynamic trials were digitised using Qualisys Track Manager (Qualisys Medical AB, Goteburg, Sweden) in order to identify anatomical and tracking markers, then exported as C3D files to Visual 3D (C-Motion, Germantown, MD, USA). All data were linearly normalized to 100% of the stance phase. GRF data and marker trajectories were smoothed with cut-off frequencies of 50 Hz at 12 Hz respectively, using a low-pass Butterworth 4th order zero lag filter. Lower extremity kinematics were quantified using an XYZ cardan sequence of rotations (where X is flexion-extension; Y is ab-adduction and is Z is internal-external rotation). Tibial internal rotation kinematics were also calculated in accordance with Eslami et al., [37]. Finally, in order to quantify contralateral pelvic drop, the coronal plane angle of the pelvic segment was quantified with respect to the laboratory. Three-dimensional kinematic measures were extracted for statistical analysis were 1) angle at footstrike, 2) peak angle and 3) range of motion (representing the angular displacement from footstrike to peak angle). Running speed was also quantified with Visual 3D using the linear velocity of the model centre of mass in the anterior direction.

In accordance with the protocol of Addison and Lieberman, [38], an impulse-momentum modelling approach was utilized to calculate effective mass as a percentage of bodyweight (% BW), which was quantified in accordance with the below equation:

$$\text{Effective mass} = \text{vertical GRF integral} / (\Delta \text{ foot vertical velocity} + \text{gravity} * \Delta \text{ time})$$

The impact peak was defined in the maximal and traditional running shoes as the first peak in vertical GRF. The time to impact peak (Δ time) was quantified as the duration from footstrike to impact peak. For runners who lacked an impact peak, in accordance with Sinclair et al., [39] we calculated based on a mean of the other runners the relative position of the impact peak as a percentage of the stance phase and positioned the impact peak here in those who did not exhibit a discernable peak in the vertical GRF time-curve. The vertical GRF integral during the period of the impact peak was calculated using a trapezoidal function. The change in foot vertical velocity (Δ foot vertical velocity) was determined as the change in vertical foot velocity between the instances of footstrike and the impact peak [40]. The velocity of the foot was quantified using the centre of mass of the foot segment in the vertical direction, within Visual 3D [39].

The loading rate expressed as bodyweight per second (BW/s) was also extracted by obtaining the peak increase in vertical GRF between adjacent data points using the first derivative function within Visual 3D. Furthermore, the GRF impulse expressed as bodyweight per millisecond (BW·ms) during the stance phase in all three anatomical directions was calculated using a trapezoidal function. The strike index (%) was calculated as the position of the centre of pressure location at footstrike, relative to the total length of the foot [41] (Figure 2a–c). A strike index of 0–33.3 % denotes a rearfoot, 33.4–66.6% a midfoot and 66.7–100% a forefoot strike pattern. Finally, vertical limb stiffness during running was quantified using a mathematical spring-mass model [42]. Vertical limb stiffness expressed as bodyweight per metre (BW/m) was calculated from the ratio of the peak vertical GRF normalized to bodyweight (BW) to the maximum vertical compression of the leg spring which was calculated as the change in limb length from footstrike to minimum length during the stance phase [43]. Limb length (m) was quantified as the vertical height of the proximal end of the thigh segment within Visual 3D.

@@@FIGURE 2 NEAR HERE@@@

Following this, data during the stance phase were exported from Visual 3D into OpenSim 3.3 software (Simtk.org). A validated musculoskeletal model was used to process the biomechanical data, which was scaled to account for the anthropometrics of each runner. The model with 12 segments, 19 degrees of freedom and 92 musculotendon actuators [44] was used to estimate lower extremity muscle and joint forces. A residual reduction algorithm [22] was firstly used to resolve dynamic inconsistency between the measured GRF's and the model's kinematics and muscle kinetics were then quantified using static optimization in accordance with Steele et al. [45].

From the above static optimization processes, the peak model muscle forces during the stance phase were extracted after being normalized to bodyweight (BW). Furthermore, the impulses (BW·ms) during the stance phases for the aforementioned muscles were also extracted using a trapezoidal function. In addition, the biceps femoris long-head, biceps femoris short-head, semitendinosus and semimembranosus muscle forces were summed to create the total hamstring muscle force and the rectus femoris, vastus lateralis, vastus medialis and vastus intermedius forces were also summed to create the total quadriceps muscle force. The peak total hamstring and quadriceps forces as well as their impulses during the stance phase were also extracted. In accordance with Besier et al., [23], the mean (%) contributions of each of the individual hamstring and quadriceps muscles to the total hamstring and total quadriceps forces during the stance phase was quantified. Finally, the mean co-contraction ratio was calculated by dividing the total hamstring force impulse by the total quadriceps impulse and peak co-contraction ratio by dividing the total hamstring force at the instance of peak patellofemoral force by the total quadriceps force at the same time point.

As muscle forces are the main determinant of joint compressive forces [46], following the static optimization process the peak compressive patellofemoral joint force (BW) was calculated via the joint reaction analyses function using the muscle forces generated from the static optimization process as inputs [22]. Furthermore, peak patellofemoral stress expressed as kilopascals per bodyweight (KPa/BW) was quantified by dividing the patellofemoral force by the joint contact area. Patellofemoral contact areas were obtained by fitting a polynomial curve to the sex specific data of Besier et al. [47], who estimated patellofemoral contact areas as a function of the knee flexion angle using magnetic resonance imaging. In addition, peak patellofemoral force (BW·ms) and stress (expressed as kilopascals per bodyweight per millisecond - KPa/BW·ms) impulses during the stance phase were extracted, also using a trapezoidal function. Finally, patellofemoral force/ stress load rates (BW/s and KPa/BW/s i.e. kilopascals per bodyweight per second) were extracted by obtaining the maximum increase in force/stress between adjacent data points using a first derivative function.

Further to the above, the patellofemoral force (BW) and patellofemoral stress (KPa/BW) experienced per mile were also quantified to explore the effect of step length on loading at this joint [48]. The number of steps needed to complete 1 mile was quantified using the step length (m), which was determined by taking the difference in the horizontal position of the foot centre of mass between the right and left limbs at foot strike from each running trial [49]. To obtain the force/ stress experienced per mile, impulse values for these measures outlined above, were multiplied by the number of steps required to run a mile.

2.4 Statistical analyses

For each outcome variable and group, means and standard deviations (SD) were calculated. Differences between four pairs of group comparisons were undertaken. 1. Overall PFP vs. Control groups were contrasted which involved all PFP runners compared against all of the asymptomatic runners. Following this subset analyses of 2. Male PFP vs. Male Control, 3. Female PFP vs. Female Control and 4. Male PFP vs. Female PFP were undertaken. To compare participant characteristics (age, mass, stature, BMI, running speed, step length, mean weekly running volume), linear mixed models with group modelled as a fixed factor and random intercepts by participants were adopted to compare pairs 1-4. Symptom duration and KOOS-PF were comparable in pair 4 only.

To compare biomechanical outcomes across the four pairs of groups, the same approach was adopted only adjusting for running speed which was modelled as continuous fixed covariate. In addition, for group comparisons from pairs 1-3 outlined above, binary logistic regression analysis was conducted with the biomechanical parameters as the predictor variables and classification into either the PFP group as the outcome variable. Finally, two-way Pearson chi-square (X^2) tests of independence with probability values calculated by Monte Carlo simulation, were used to assess differences in the distribution of the three-foot strike patterns between the pairs. Only, parameters identified to be significantly different between control and PFP groups using linear mixed model analyses, were considered for the regression analyses. All statistical analyses were conducted using SPSS v27 (IBM, SPSS). For linear mixed models, the mean difference (b), t -value and 95% confidence intervals of the difference are presented. For binary logistic regression analyses, the odds ratio (OR) and 95% confidence intervals of the odds ratio are presented. Statistical significance for all analyses was accepted at the $P < 0.05$ level. In the interests of conciseness and clarity only variables that presented with statistical significance are presented in the results section.

3. Results

3.1 Participant characteristics

Comparisons between all groups for (age, BMI, running speed, mean weekly running volume, symptom duration and KOOS-PF) were non-significant ($p = 0.12$ – 0.97). However, mass ($b = 15.98$, (95% CI = $7.25 - 24.69$), $t = 3.71$, $p < 0.001$) and stature ($b = 0.11$, (95% CI = $0.08 - 0.15$), $t = 5.68$, $p < 0.001$) were significantly greater in symptomatic males compared to females with PFP. Furthermore, step length was shown to be significantly greater in the control group compared to overall PFP ($b = 0.27$, (95% CI = $0.14 - 0.40$), $t = 4.023$, $p < 0.001$) and in male controls in relation to male PFP runners ($b = 0.15$, (95% CI = $0.07 - 0.22$), $t = 4.067$, $p < 0.001$) (Table 1).

@@@TABLE 1 NEAR HERE@@@

3.2 Overall PFP vs. Control

Knee flexion at footstrike, knee external rotation at footstrike, peak knee flexion, knee coronal plane ROM, knee transverse plane ROM and ankle transverse plane ROM were significantly larger in the control group. Hip external rotation at footstrike, hip sagittal plane ROM, knee abduction at footstrike, ankle eversion at footstrike, ankle external rotation at

footstrike, peak ankle eversion and pelvic ROM were significantly larger in the PFP group (Table 2).

@@@TABLE 2 NEAR HERE@@@

In addition, the strike index was significantly greater in the control group. The patellofemoral force per mile and patellofemoral stress per mile were significantly larger in the PFP group (Table 3).

@@@TABLE 3 NEAR HERE@@@

Peak adductor longus force, flexor digitorum longus impulse, flexor hallucis longus impulse, gluteus minimus impulse, peroneus tertius impulse, semitendinosus impulse, peak semitendinosus force, soleus impulse, semitendinosus %, vastus intermedius %, vastus lateralis %, vastus medialis and mean co-contraction ratio were significantly greater in the control group. The peak extensor hallucis longus force, peak lateral gastrocnemius force, rectus femoris impulse, peak semimembranosus force, biceps femoris short head % and rectus femoris % were significantly greater in the PFP group (Table 4).

@@@TABLE 4 NEAR HERE@@@

The control group was comprised of 25 rearfoot strikers, 11 midfoot strike runners and 4 forefoot strikers, whereas the PFP group had 37 rearfoot strike runners, 2 midfoot strikers and 1 forefoot runner. The chi-squared test was significant ($X^2_{(2)} = 10.35$, $P=0.006$) with statistically more rearfoot runners in the PFP group and more mid and forefoot strikers in the control group.

3.3 Male PFP vs. Male Control

The ankle transverse plane ROM was significantly greater in the control group. The ankle eversion at footstrike, ankle internal rotation at footstrike, peak ankle eversion, and pelvic ROM were significantly greater in the PFP group (Table 5).

@@@TABLE 5 NEAR HERE@@@

In addition, adductor magnus impulse, gluteus minimus impulse, peroneus brevis impulse, peak peroneus brevis force, semitendinosus impulse, peak semitendinosus force, peak soleus force, semitendinosus %, vastus intermedius %, vastus lateralis % and vastus medialis % were significantly greater in the control group. Rectus femoris % was however significantly greater in the PFP group (Table 6).

@@@TABLE 6 NEAR HERE@@@

The control group was comprised of 19 rearfoot strikers, 4 midfoot strike runners and 2 forefoot strikers, whereas the male PFP group had 22 rearfoot strike runners, 2 midfoot strikers and 1 forefoot runner. The chi-squared test was non-significant ($X^2_{(2)} = 1.56$, $P=0.458$).

3.4 Female PFP vs. Female Control

Knee external rotation at footstrike and knee transverse plane ROM were significantly greater in the control group. Knee abduction at footstrike, ankle dorsiflexion at footstrike, ankle eversion at footstrike, ankle external rotation at footstrike, peak ankle dorsiflexion, peak ankle eversion and pelvic ROM were significantly larger in the PFP group (Table 7).

@@@TABLE 7 NEAR HERE@@@

In addition, the strike index was significantly greater in the control group. The vertical GRF impulse, patellofemoral force impulse, peak patellofemoral force, peak patellofemoral stress, patellofemoral stress loading rate, patellofemoral stress impulse and patellofemoral stress per mile were significantly larger in the PFP group (Table 8).

@@@TABLE 8 NEAR HERE@@@

Peak adductor longus force Semitendinosus %, vastus intermedius %, vastus lateralis % and vastus medialis % were significantly greater in the control group. Peak biceps femoris short head force, peak extensor hallucis longus force, peak lateral gastrocnemius force, peak medial gastrocnemius force, peak peroneus longus force, psoas impulse, rectus femoris impulse, peak semimembranosus force, total quadriceps impulse, peak total quadriceps force, peak vastus intermedius force, vastus intermedius impulse, peak vastus medialis force, vastus medialis impulse and rectus femoris % were significantly greater in the PFP group (Table 9).

@@@TABLE 9 NEAR HERE@@@

The control group was comprised of 7 rearfoot strikers, 6 midfoot strike runners and 2 forefoot strikers, whereas the female PFP group had 15 rearfoot strike runners. The chi-squared test was significant ($X^2_{(2)} = 11.05$, $P=0.004$) with statistically more rearfoot runners in the female PFP group and more mid and forefoot strikers in the control group.

3.5 Male PFP vs. Female PFP

Hip flexion at footstrike, peak hip adduction, knee flexion at footstrike and peak knee flexion were significantly larger in the female PFP group. Knee transverse plane ROM was significantly greater in the male PFP group (Table 10).

@@@TABLE 10 NEAR HERE@@@

The loading rate was significantly larger in the female PFP group (Table 11).

@@@TABLE 11 NEAR HERE@@@

Biceps femoris long head impulse, peak biceps femoris short head force, gluteus maximus impulse, peak gluteus maximus force, gluteus medius impulse, peak gluteus medius force, gluteus minimus impulse, peak gluteus minimus force, peak gracilis force, peak iliopsoas force, peak sartorius force, semimembranosus impulse, peak semimembranosus force, semitendinosus impulse, peak semitendinosus force, tensor fasciae latae impulse, peak tensor fasciae latae force, tibialis anterior impulse, total hamstring impulse, peak total hamstring force vastus intermedius %, vastus lateralis %, mean co-contraction ratio and peak co-contraction ratio were significantly greater in the female PFP group. Extensor digitorum longus impulse, flexor digitorum longus impulse, rectus femoris impulse and rectus femoris % were significantly greater in the male PFP group (Table 12).

@@@TABLE 12 NEAR HERE@@@

The male PFP group was comprised of 2 rearfoot strikers, 1 midfoot strike runners and 1 forefoot strikers, whereas the female PFP group had 15 rearfoot strike runners. The chi-squared test was non-significant ($X^2_{(2)} = 1.95$, $P=0.378$).

4. Discussion

Using a retrospective case-control approach, this study evaluated running biomechanics in patients with and without PFP via a musculoskeletal simulation-based approach, also investigated potential distinctions in PFP presentation between the sexes. This study represents the first exploration of the biomechanical parameters linked to the aetiology of

PFP using musculoskeletal simulation, and may yield important clinical information towards the cessation of symptom initiation and the improvement in PFP management.

Notably, patellofemoral stress indices were statistically greater in the PFP group compared to controls. This supports the notion that in runners, PFP is mediated by increased joint loading, and concurs with the findings of Brechter & Powers, [50], who found enhanced patellofemoral joint stress in those with PFP using a mixed sample. The subset analyses from this study importantly revealed that enhanced patellofemoral joint stress in PFP runners was driven by females, as no differences were evident in males. The female PFP group exhibited more pronounced and coherent differences in patellofemoral joint loading, importantly highlighting potential sex differences in PFP aetiology. This emphasizes the need for different treatment approaches in male and female runners and management strategies seeking to reduce patellofemoral joint stress are likely to be most efficacious in female runners. Patellofemoral joint loading is primarily actuated by forces generated within the quadriceps [51]. Therefore, enhanced patellofemoral joint stress in the female PFP group compared, was mediated as a function of the corresponding increases in forces which were observed consistently across all four quadriceps muscles.

A significantly more anterior footstrike position was shown in the control group in relation to the PFP group, which when contextualized via the strike index, revealed a rearfoot strike in the PFP group and a midfoot pattern in controls. The subset analyses again showed that this was mediated by female runner's and the chi-square analyses support this, as significantly more rearfoot strikers were shown in the overall and female PFP groups. Daoud et al., [52] showed that runners who habitually adopt a rearfoot strike pattern are more than twice as likely to experience a chronic running injury. Crucially, this investigation provides further precision to this observation, as in female runners the utilization of a rearfoot strike pattern makes runners significantly more likely to experience PFP. It is likely that the increased quadriceps muscle forces and patellofemoral loading shown overall and in the female PFP group, was mediated via the rearfoot strike pattern adopted in relation to control runners. Utilization of a rearfoot strike pattern places increased demands on the knee extensors [53], and patellofemoral stress is greater in rearfoot strikers [54]. From a clinical perspective, transitioning from a rearfoot strike pattern has been shown to attenuate patellofemoral joint loading and improve pain symptoms in male runners, although no such intervention has been undertaken in females [25]. Therefore, gait-retraining approaches which allow female PFP runners to modify their habitual footstrike pattern, may be a particularly important clinical intervention.

From a kinematic and temporal perspective, peak knee flexion and step length were lower in the PFP group and knee flexion lower in male symptomatic runners compared to females with PFP. Individuals with PFP utilize reduced knee flexion [55, 56], which allied to the reductions in step length, may represent an attempt to minimize pain symptoms by mediating patellofemoral stress [49, 55]. Reduced knee flexion in both the overall PFP and male PFP runners, may be indicative of kinesiophobia, a renowned complaint in patients with PFP [10, 56]. Kinesiophobia, is more severe in males with chronic musculoskeletal pain [57, 58]. Therefore, as knee bracing [59] and exercise therapy [60] have been shown to improve kinesiophobia in PFP, this suggests there is scope for targeted interventions towards male symptomatic runners.

The overall and male and female subset comparisons showed that peak eversion was significantly greater in PFP runners. Rearfoot eversion has been linked to the aetiology of PFP by theoretical modelling analyses [61], and Dierks et al., [18] showed significantly larger eversion in PFP runners in a mixed sample, though this investigation is the first to show increased eversion in sex specific subset analyses. From a clinical standpoint, it appears that strategies centred around reducing rearfoot eversion are important to the treatment of PFP in both male and female runners. Conservative modalities such as foot orthoses and ankle braces have been shown to mediate statistical reductions in peak ankle eversion in both male and female runners [62, 63]. No analyses have been undertaken

using ankle bracing, however previous intervention studies have shown foot orthoses to be effective in treating PFP with greater rearfoot eversion angles at baseline predicting orthotic efficacy [64, 65].

This investigation also revealed in the overall and male and female subset comparisons, that contralateral pelvic drop ROM was significantly larger in PFP runners. This supports Bramah et al., [66] who showed that injured runners demonstrated greater contralateral pelvic drop and Willson & Davis, [19] who indicated that runners with PFP exhibited significantly greater contralateral pelvic drop. Enhanced contralateral pelvic drop negatively influences patellofemoral joint biomechanics by elongating the ipsilateral iliotibial band, generating a greater laterally directed force on the patella [67, 68]. Increased contralateral pelvic drop during weightbearing activities and in PFP patients is considered a clinical sign of diminished strength or neuromuscular function at the hip [66], which results in diminished capacity to stabilize the pelvis. This investigation supports this notion, as forces in the muscles primarily involved in stabilizing the pelvis were found to be statistically greater in the control group across all of the aforementioned group comparisons. Importantly, real-time gait retraining has been shown to significantly attenuate contralateral pelvic drop and pain symptoms in runners with PFP and that hip strengthening is able to mediate statistical improvements in PFP pain symptoms [69]. The findings from the current investigation importantly support the collective utilization of these intervention modalities in runners of both sexes.

The overall as well as the male and female comparisons showed that % contributions of the rectus femoris were consistently greater in PFP runners, and that the same was true in symptomatic male runners. This suggests that over utilization of the rectus femoris is a potentially important biomechanical mechanism in PFP, and able to differentiate between sexes in symptomatic runners. Although, the mechanistic influence of this finding relating to the aetiology of PFP is not known, it is nonetheless a potentially clinically meaningful neuromuscular observation that may aid in the management of PFP in runners of both sexes. Furthermore, the overall and subgroup comparison of symptomatic runners showed an enhanced co-contraction ratio in control runners and symptomatic females. This observation opposes those of Besier et al., [23], showing no differences between PFP and healthy controls during running, although this investigation utilized electromyography rather than muscle forces. Nonetheless, it is proposed that these observations are mediated as a function of the enhanced hamstring and reduced rectus femoris muscle forces that were evident in the above overall and subgroup comparisons. It remains unknown whether these alterations in muscle kinetics are adaptive in response to pain or if they are causative, but it can be speculated that increased co-contraction might enhance knee joint stability and benefit the position of the patella within the trochlear groove [70].

In contrary to the observations of both Willy et al., [20] and Neal et al., [4] our findings showed that females with PFP did not exhibit a greater peak hip adduction angle compared to controls. However, in agreement with Willy et al., [20] our results show that female runners with PFP ran with statistically larger hip adduction in relation to symptomatic males. Previous intervention trials show that targeted gait-retraining reduces peak hip adduction during running and improve pain and function in runners with PFP [71]. The observations from this investigation suggest that these interventions are likely to be most effective in females and the implementation of gait re-training to attenuate hip adduction in symptomatic female runners is encouraged. Taking into account the range of sex specific differences in running biomechanics between symptomatic runners, demonstrates sex specificity in terms of PFP presentation. Indicating that future clinical investigations whether, should present aetiological data for both sexes, and that treatment modalities should also be correspondingly sex specific.

Like any investigation, there are limitations to the current study, and the findings should be contextualized in line with these limitations. Firstly, the retrospective nature of the study design means that it is not possible to determine whether the biomechanical

differences between symptomatic runners and healthy controls are adaptations in the presence of anterior knee pain or if they are causative parameters. Owing to the expense and difficulty of undertaking analyses of this nature, there are a lack of prospective analyses identifying risk factors for PFP, so this is a clear avenue for future research. In addition, PFP is a multifactorial condition; therefore, although the current investigation explored a plethora of parameters using musculoskeletal simulation, it was beyond the scope of this study to determine the non-biomechanical risk factors that may be important to the aetiology of PFP. Future analyses should seek to investigate additional measurements that may be important to the aetiology and clinical management of PFP.

5. Conclusions

This investigation augments the existing literature in clinical biomechanics by providing a retrospective exploration of running biomechanics in runners with and without PFP using musculoskeletal simulation. With overall PFP as well as female PFP runners exhibiting enhanced patellofemoral stress in addition to a significantly lower strike index, peak eversion and contralateral pelvic drop being greater in overall, male and female PFP runners, symptomatic females exhibiting a significantly larger peak hip adduction compared to PFP males and peak knee flexion being reduced in overall PFP and male PFP runners compared to symptomatic females; this investigation highlights potential risk factors for PFP that may be addressed through focused intervention modalities. Specifically, gait re-training approaches allowing female PFP runners to modify their habitual footstrike pattern and attenuate hip joint adduction, foot orthoses to reduce eversion and targeted gait re-training to reduce pelvic drop in male and female PFP runners and knee bracing and exercise therapy to mediate potential symptoms of kinesiophobia in male PFP runners, appear to represent particularly important clinical interventions.

Author Contributions: Conceptualization: JKS & PJT, methodology: JKS, NC & PJT, processing: JKS & NC, data analysis: JKS, writing JKS, NC & PJT. All authors have read and agreed to the published version of the manuscript.

Institutional Review Board Statement: Participants provided written informed consent and ethical approval (REF: STEMH 424) was obtained from the University of Central Lancashire, in accordance with the principles documented in the Declaration of Helsinki.

Informed Consent Statement: Participants provided written informed consent.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Lee, D. C., Pate, R. R., Lavie, C. J., Sui, X., Church, T. S., & Blair, S. N. (2014). Leisure-time running reduces all-cause and cardiovascular mortality risk. *Journal of the American College of Cardiology*, 64(5), 472-481.
- Taunton, J. E., Ryan, M. B., Clement, D. B., McKenzie, D. C., Lloyd-Smith, D. R., & Zumbo, B. D. (2002). A retrospective case-control analysis of 2002 running injuries. *British Journal of Sports Medicine*, 36(2), 95-101.
- Aicale, R., Tarantino, D., & Maffulli, N. (2018). Overuse injuries in sport: a comprehensive overview. *Journal of Orthopaedic Surgery and Research*, 13(1), 1-11.
- Neal, B. S., Barton, C. J., Birn-Jeffery, A., & Morrissey, D. (2019). Increased hip adduction during running is associated with patellofemoral pain and differs between males and females: A case-control study. *Journal of Biomechanics*, 91, 133-139.
- Thijs, Y., Pattyn, E., Van Tiggelen, D., Rombaut, L., & Witvrouw, E. (2011). Is hip muscle weakness a predisposing factor for patellofemoral pain in female novice runners? A prospective study. *The American Journal of Sports Medicine*, 39(9), 1877-1882.
- Robinson, R. L., & Nee, R. J. (2007). Analysis of hip strength in females seeking physical therapy treatment for unilateral patellofemoral pain syndrome. *Journal of Orthopaedic & Sports Physical Therapy*, 37(5), 232-238.
- Nimon, G., Murray, D., Sandow, M., & Goodfellow, J. (1998). Natural history of anterior knee pain: a 14-to 20-year follow-up of nonoperative management. *Journal of Pediatric Orthopaedics*, 18(1), 118-122.
- Thomas, M. J., Wood, L., Selfe, J., & Peat, G. (2010). Anterior knee pain in younger adults as a precursor to subsequent patellofemoral osteoarthritis: a systematic review. *BMC Musculoskeletal Disorders*, 11(1), 1-8.
- Blond, L., & Hansen, L. (1998). Patellofemoral pain syndrome in athletes: a 5.7-year retrospective follow-up study of 250 athletes. *Acta Orthopaedica Belgica*, 64(4), 393-400.
- Maclachlan, L. R., Matthews, M., Hodges, P. W., Collins, N. J., & Vicenzino, B. (2018). The psychological features of patellofemoral pain: a cross-sectional study. *Scandinavian Journal of Pain*, 18(2), 261-271.

11. Lack, S., Neal, B., Silva, D. D. O., & Barton, C. (2018). How to manage patellofemoral pain—Understanding the multifactorial nature and treatment options. *Physical Therapy in Sport*, 32, 155-166. 557
12. Neal, B., Lack, S., Lankhorst, N., Raye, A., Morrissey, D., & Van Middelkoop, M. (2019). 7 Risk factors for patellofemoral pain: a systematic review & meta-analysis. *British Journal of Sports Medicine*, 53(5), 270-281. 558
13. Noehren, B., Hamill, J., & Davis, I. (2013). Prospective evidence for a hip etiology in patellofemoral pain. *Medicine and Science in Sports and Exercise*, 45(6), 1120-1124. 559
14. Fox, A., Ferber, R., Saunders, N., Osis, S., & Bonacci, J. (2018). Gait kinematics in individuals with acute and chronic patellofemoral pain. *Medicine and Science in Sports and Exercise*, 50, 502-509. 560
15. Noehren, B., Pohl, M. B., Sanchez, Z., Cunningham, T., & Lattermann, C. (2012). Proximal and distal kinematics in female runners with patellofemoral pain. *Clinical biomechanics*, 27(4), 366-371. 561
16. Noehren, B., Sanchez, Z., Cunningham, T., & McKeon, P. O. (2012). The effect of pain on hip and knee kinematics during running in females with chronic patellofemoral pain. *Gait & Posture*, 36(3), 596-599. 562
17. Esculier, J. F., Roy, J. S., & Bouyer, L. J. (2015). Lower limb control and strength in runners with and without patellofemoral pain syndrome. *Gait & posture*, 41(3), 813-819. 563
18. Dierks, T. A., Manal, K. T., Hamill, J., & Davis, I. S. (2008). Proximal and distal influences on hip and knee kinematics in runners with patellofemoral pain during a prolonged run. *Journal of Orthopaedic & Sports Physical Therapy*, 38(8), 448-456. 564
19. Willson, J. D., & Davis, I. S. (2008). Lower extremity mechanics of females with and without patellofemoral pain across activities with progressively greater task demands. *Clinical Biomechanics*, 23(2), 203-211. 565
20. Willy, R. W., Manal, K. T., Witvrouw, E. E., & Davis, I. S. (2012). Are mechanics different between male and female runners with patellofemoral pain?. *Medicine and Science in Sports and Exercise*, 44(11), 2165-2171. 566
21. Pereiro-Buceta, H., Becerro-de-Bengoa-Vallejo, R., Losa-Iglesias, M. E., López-López, D., Navarro-Flores, E., Martínez-Jiménez, E. M., & Calvo-Lobo, C. (2021). The Effect of Simulated Leg-Length Discrepancy on the Dynamic Parameters of the Feet during Gait. *Healthcare*, 9, (8), 932. 567
22. Delp, S. L., Anderson, F. C., Arnold, A. S., Loan, P., Habib, A., John, C. T., ... & Thelen, D. G. (2007). OpenSim: open-source software to create and analyze dynamic simulations of movement. *IEEE transactions on biomedical engineering*, 54(11), 1940-1950. 568
23. Besier, T. F., Fredericson, M., Gold, G. E., Beaupré, G. S., & Delp, S. L. (2009). Knee muscle forces during walking and running in patellofemoral pain patients and pain-free controls. *Journal of Biomechanics*, 42(7), 898-905. 569
24. Rosner, B. (2015). *Fundamentals of biostatistics*. Cengage learning. 570
25. Sinclair, J. K. (2016). Effects of a 10-week footstrike transition in habitual rearfoot runners with patellofemoral pain. *Comparative Exercise Physiology*, 12(3), 141-150. 571
26. Sinclair, J. (2014). Effects of barefoot and barefoot inspired footwear on knee and ankle loading during running. *Clinical Biomechanics*, 29(4), 395-399. 572
27. Sinclair, J., Janssen, J., Richards, J. D., Butters, B., Taylor, P. J., & Hobbs, S. J. (2018). Effects of a 4-week intervention using semi-custom insoles on perceived pain and patellofemoral loading in targeted subgroups of recreational runners with patellofemoral pain. *Physical Therapy in Sport*, 34, 21-27. 573
28. Sinclair, J., & Selfe, J. (2015). Sex differences in knee loading in recreational runners. *Journal of Biomechanics*, 48(10), 2171-2175. 574
29. Crossley, K. M., van Middelkoop, M., Callaghan, M. J., Collins, N. J., Rathleff, M. S., & Barton, C. J. (2016). 2016 Patellofemoral pain consensus statement from the 4th International Patellofemoral Pain Research Retreat, Manchester. Part 2: recommended physical interventions (exercise, taping, bracing, foot orthoses and combined interventions). *British Journal of Sports Medicine*, 50(14), 844-852. 575
30. Crossley, K. M., Macri, E. M., Cowan, S. M., Collins, N. J., & Roos, E. M. (2018). The patellofemoral pain and osteoarthritis subscale of the KOOS (KOOS-PF): development and validation using the COSMIN checklist. *British Journal of Sports Medicine*, 52(17), 1130-1136. 576
31. Sinclair, J. K., Edmundson, C. J., Brooks, D., & Hobbs, S. J. (2011). Evaluation of kinematic methods of identifying gait Events during running. *International Journal of Sports Science and Engineering*, 5(3), 188-192. 577
32. Cappozzo, A., Catani, F., Della Croce, U., & Leardini, A. (1995). Position and orientation in space of bones during movement: anatomical frame definition and determination. *Clinical Biomechanics*, 10(4), 171-178. 578
33. Sinclair, J., Hebron, J., & Taylor, P. J. (2014). The influence of tester experience on the reliability of 3D kinematic information during running. *Gait & Posture*, 40(4), 707-711. 579
34. Graydon, R., Fewtrell, D. J., Atkins, S. J., & Sinclair, J. K. (2015). The test-retest reliability of different ankle joint center location techniques. *The Foot and Ankle Online Journal*, 8(1), 22-26. 580
35. Sinclair, J., Hebron, J., & Taylor, P. J. (2015). The test-retest reliability of knee joint center location techniques. *Journal of Applied Biomechanics*, 31(2), 117-121. 581
36. Sinclair, J., Taylor, P. J., Currigan, G., & Hobbs, S. J. (2014). The test-retest reliability of three different hip joint centre location techniques. *Movement & Sport Sciences-Science & Motricité*, (83), 31-39. 582
37. Eslami, M., Begon, M., Farahpour, N., & Allard, P. (2007). Forefoot–rearfoot coupling patterns and tibial internal rotation during stance phase of barefoot versus shod running. *Clinical Biomechanics*, 22(1), 74-80. 583
38. Addison, B. J., & Lieberman, D. E. (2015). Tradeoffs between impact loading rate, vertical impulse and effective mass for walkers and heel strike runners wearing footwear of varying stiffness. *Journal of Biomechanics*, 48(7), 1318-1324. 584

39. Sinclair, J., Stainton, P., & Hobbs, S. J. (2018). Effects of barefoot and minimally shod footwear on effective mass—Implications for transient musculoskeletal loading. *Kinesiology*, 50(2), 165–171.
40. Chi, K. J., & Schmitt, D. (2005). Mechanical energy and effective foot mass during impact loading of walking and running. *Journal of Biomechanics*, 38(7), 1387–1395.
41. Squadrone, R., Rodano, R., Hamill, J., & Preatoni, E. (2015). Acute effect of different minimalist shoes on foot strike pattern and kinematics in rearfoot strikers during running. *Journal of Sports Sciences*, 33(11), 1196–1204.
42. Blickhan, R. (1989). The spring-mass model for running and hopping. *Journal of Biomechanics*, 22(11–12), 1217–1227.
43. Farley, C. T., & Morgenroth, D. C. (1999). Leg stiffness primarily depends on ankle stiffness during human hopping. *Journal of Biomechanics*, 32(3), 267–273.
44. Lerner, Z. F., DeMers, M. S., Delp, S. L., & Browning, R. C. (2015). How tibiofemoral alignment and contact locations affect predictions of medial and lateral tibiofemoral contact forces. *Journal of Biomechanics*, 48(4), 644–650.
45. Steele, K. M., DeMers, M. S., Schwartz, M. H., & Delp, S. L. (2012). Compressive tibiofemoral force during crouch gait. *Gait & Posture*, 35(4), 556–560.
46. Herzog, W., Longino, D., & Clark, A. (2003). The role of muscles in joint adaptation and degeneration. *Langenbeck's Archives of Surgery*, 388(5), 305–315.
47. Besier, T. F., Draper, C. E., Gold, G. E., Beaupré, G. S., & Delp, S. L. (2005). Patellofemoral joint contact area increases with knee flexion and weight-bearing. *Journal of Orthopaedic Research*, 23(2), 345–350.
48. Sinclair, J., Taylor, P. J., & Liles, N. B. (2020). Effects of running with minimal and conventional footwear in habitual and non-habitual users: a musculoskeletal simulation and statistical parametric mapping based approach. *Footwear Science*, 12(1), 25–38.
49. Sinclair, J., Richards, J., Selfe, J., Fau-Goodwin, J., & Shore, H. (2016). The influence of minimalist and maximalist footwear on patellofemoral kinetics during running. *Journal of Applied Biomechanics*, 32(4), 359–364.
50. Brechter, H., & Powers, C. M. (2002). Patellofemoral stress during walking in persons with and without patellofemoral pain. *Medicine and Science in Sports and Exercise*, 34(10), 1582–1593.
51. Loudon, J. K. (2016). Biomechanics and pathomechanics of the patellofemoral joint. *International Journal of Sports Physical Therapy*, 11(6), 820–830.
52. Daoud, A. I., Geissler, G. J., Wang, F., Saretsky, J., Daoud, Y. A., & Lieberman, D. E. (2012). Foot strike and injury rates in endurance runners: a retrospective study. *Medicine and Science in Sports and Exercise*, 44(7), 1325–1334.
53. Powers, C. M., Ho, K. Y., Chen, Y. J., Souza, R. B., & Farrokhi, S. (2014). Patellofemoral joint stress during weight-bearing and non—weight-bearing quadriceps exercises. *Journal of Orthopaedic & Sports Physical Therapy*, 44(5), 320–327.
54. Kulmala, J. P., Avela, J. A. N. N. E., Pasanen, K. A. T. I., & Parkkari, J. A. R. I. (2013). Forefoot strikers exhibit lower running-induced knee loading than rearfoot strikers. *Medicine and Science in Sports and Exercise*, 45(12), 2306–2313.
55. Crossley, K. M., Cowan, S. M., Bennell, K. L., & McConnell, J. (2004). Knee flexion during stair ambulation is altered in individuals with patellofemoral pain. *Journal of Orthopaedic Research*, 22(2), 267–274.
56. de Oliveira Silva, D., Barton, C. J., Briani, R. V., Taborda, B., Ferreira, A. S., Pazzinatto, M. F., & de Azevedo, F. M. (2019). Kinesiophobia, but not strength is associated with altered movement in women with patellofemoral pain. *Gait & Posture*, 68, 1–5.
57. Bränström, H., & Fahlström, M. (2008). Kinesiophobia in patients with chronic musculoskeletal pain: differences between men and women. *Journal of Rehabilitation Medicine*, 40(5), 375–380.
58. Rovner, G. S., Sunnerhagen, K. S., Björkdahl, A., Gerdle, B., Börsbo, B., Johansson, F., & Gillanders, D. (2017). Chronic pain and sex-differences; women accept and move, while men feel blue. *PloS one*, 12(4), e0175737.
59. Priore, L. B., Lack, S., Garcia, C., Azevedo, F. M., & de Oliveira Silva, D. (2020). Two weeks of wearing a knee brace compared with minimal intervention on kinesiophobia at 2 and 6 weeks in people with patellofemoral pain: a randomized controlled trial. *Archives of Physical Medicine and Rehabilitation*, 101(4), 613–623.
60. Smith, B. E., Hendrick, P., Bateman, M., Moffatt, F., Rathleff, M. S., Selfe, J., & Logan, P. (2019). A loaded self-managed exercise programme for patellofemoral pain: a mixed methods feasibility study. *BMC Musculoskeletal Disorders*, 20(1), 1–13.
61. Tiberio, D. (1987). The effect of excessive subtalar joint pronation on patellofemoral mechanics: a theoretical model. *Journal of orthopaedic & Sports Physical Therapy*, 9(4), 160–165.
62. Sinclair, J., Ingram, J., Taylor, P. J., & Chockalingam, N. (2019). Acute effects of different orthoses on lower extremity kinetics and kinematics during running; a musculoskeletal simulation analysis. *Acta of Bioengineering and Biomechanics*, 21(4).
63. Graydon, R., Fewtrell, D., Atkins, S., & Sinclair, J. (2017). The effects of ankle protectors on lower limb kinematics in male football players: a comparison to braced and unbraced ankles. *Comparative Exercise Physiology*, 13(4), 251–258.
64. Barton, C. J., Levinger, P., Webster, K. E., & Menz, H. B. (2011). Kinematics associated with foot pronation in individuals with patellofemoral pain syndrome: a case-control study. *Journal of Foot and Ankle Research*, 4(1), 1–1.
65. Munuera, P. V., & Mazoterias-Pardo, R. (2011). Benefits of custom-made foot orthoses in treating patellofemoral pain. *Prosthetics and Orthotics International*, 35(4), 342–349.
66. Bramah, C., Preece, S. J., Gill, N., & Herrington, L. (2018). Is there a pathological gait associated with common soft tissue running injuries?. *The American Journal of Sports Medicine*, 46(12), 3023–3031.
67. Puniello, M. S. (1993). Iliotibial band tightness and medial patellar glide in patients with patellofemoral dysfunction. *Journal of Orthopaedic & Sports Physical Therapy*, 17(3), 144–148.

68. Wu, C. C., & Shih, C. H. (2004). The influence of iliotibial tract on patellar tracking. *Orthopedics*, 27(2):199-203.
69. Santos, T. R., Oliveira, B. A., Ocarino, J. M., Holt, K. G., & Fonseca, S. T. (2015). Effectiveness of hip muscle strengthening in patellofemoral pain syndrome patients: a systematic review. *Brazilian Journal of Physical Therapy*, 19, 167-176.
70. Farahmand, F., Sejiavongse, W., & Amis, A. A. (1998). Quantitative study of the quadriceps muscles and trochlear groove geometry related to instability of the patellofemoral joint. *Journal of Orthopaedic Research*, 16(1), 136-143.
71. Willy, R. W., Scholz, J. P., & Davis, I. S. (2012). Mirror gait retraining for the treatment of patellofemoral pain in female runners. *Clinical Biomechanics*, 27(10), 1045-1051.




Figure 1: a. Experimental marker locations and b. trunk, pelvis, thigh, shank and foot segments, with segment co-ordinate system axes (R = right & L = left), (TR = trunk, P = pelvis, T = thigh, S = shank & F = foot), (X = sagittal, Y = coronal & Z = transverse planes).




Figure 2: Definitions of foot strike modalities according to the strike index [39]. a. Rear-foot strike; b. mid-foot strike & c. fore-foot strike.

691
692

Table 1: Participant characteristics.

	Control		PFP		Male control		Male PFP		Female control		Female PFP	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Age (yrs)	33.25	10.78	35.85	9.29	33.87	10.42	36.60	9.07	32.76	11.55	34.60	9.83
Mass (kg)	70.91	9.65	73.01	15.19	75.65	9.81	80.00	14.59	65.21	8.31	64.02	10.34
Stature (m)	1.73	0.09	1.73	0.08	1.74	0.07	1.77	0.07	1.68	0.10	1.66	0.04
BMI (kg/m ²)	24.55	3.26	24.46	3.36	23.79	3.00	25.39	3.44	23.22	3.66	23.20	2.83
Speed (m/s)	3.89	0.43	3.66	0.58	3.90	0.52	3.72	0.60	3.86	0.27	3.56	0.55
Step length (m)	1.47	0.36	1.20	0.23	1.50	0.29	1.21	0.23	1.43	0.45	1.19	0.24
Mean weekly running volume (km)	27.12	15.51	30.09	19.62	29.04	19.14	30.94	19.72	25.21	10.96	29.24	19.99
Symptom duration (Months)	N/A		48.73	32.06	N/A		42.32	26.84	N/A		55.13	36.10
KOOS-PF	N/A		61.25	15.37	N/A		58.10	15.34	N/A		66.50	14.40

Table 2: Three-dimensional kinematics in overall PFP participants and controls.

		Control		Patellofemoral		<i>t</i>	<i>b</i>	<i>b</i> (95% CI)	<i>p</i>	OR	<i>OR</i> (95% CI)
		Mean	SD	Mean	SD						
Sagittal plane	Hip ROM (°)	0.60	1.12	1.78	1.93	3.39	1.21	0.50 - 1.92	0.001	1.75	1.18 - 2.59
	Knee angle at footstrike (°)	16.83	8.81	11.85	5.41	3.17	5.23	1.95 - 8.51	0.002	0.90	0.83 - 0.97
	Peak knee flexion (°)	43.95	6.71	40.29	7.80	2.08	3.41	0.15 - 6.68	0.041	0.93	0.873 - 1.00
Coronal plane	Pelvic ROM (°)	1.73	1.45	3.16	2.05	3.33	1.32	0.53 - 2.12	0.001	1.54	1.15 - 2.05
	Knee angle at footstrike (°)	0.50	5.23	-2.71	4.63	2.69	2.99	0.78 - 5.20	0.009	0.88	0.80 - 0.97
	Knee ROM (°)	6.31	4.06	3.83	2.58	3.14	2.41	0.88 - 3.94	0.002	0.81	0.70 - 0.94
	Ankle angle at footstrike (°)	5.66	5.04	-4.39	6.05	8.02	10.10	7.59 - 12.60	<0.001	0.70	0.60 - 0.82
	Peak ankle eversion (°)	-8.15	4.19	-15.09	7.39	5.26	7.12	4.43 - 9.82	<0.001	0.78	0.70 - 0.88
Transverse plane	Hip angle at footstrike (°)	3.66	10.30	-1.45	9.50	2.08	4.64	0.20 - 9.07	0.041	0.95	0.91 - 1.01
	Knee angle at footstrike (°)	-9.96	8.62	-4.61	7.75	2.66	4.90	1.23 - 8.56	0.009	1.08	1.01 - 1.15
	Knee ROM (°)	17.48	6.66	12.98	4.86	3.08	3.98	1.42 - 6.56	0.003	0.88	0.80 - 0.97

	Ankle angle at footstrike (°)	-6.38	7.90	-15.68	4.74	6.37	9.38	6.45 - 12.32	<0.001	0.80	0.721 - 0.89
	Ankle ROM (°)	7.82	4.16	0.61	1.03	10.42	7.11	5.75 - 8.47	<0.001	0.35	0.21 - 0.56

Table 3: Joint loading and temporal indices in overall PFP participants and controls.

	Control		Patellofemoral		<i>t</i>	<i>b</i>	<i>b</i> (95% CI)	<i>p</i>	<i>OR</i>	<i>OR</i> (95% CI)
	Mean	SD	Mean	SD						
Strike index (%)	33.57	22.09	17.75	16.43	3.52	15.47	6.71 - 24.22	0.001	0.946	0.91 - 0.98
Step length (m)	1.47	0.36	1.20	0.23	3.65	0.24	0.11 - 0.38	<0.001	0.043	0.01 - 0.33
Patellofemoral force per mile (BW)	440.24	204.84	550.33	192.08	2.14	94.46	6.55 - 182.37	0.036	1.003	1.00 - 1.01
Patellofemoral stress per mile (KPa/ BW)	812.93	363.66	1047.49	333.02	2.64	203.77	50.07 - 357.47	0.01	1.002	1.001 - 1.004

Table 4: Muscle force indices in overall PFP participants and controls.

	Control		Patellofemoral		<i>t</i>	<i>b</i>	<i>b</i> (95% CI)	<i>p</i>	<i>OR</i>	<i>OR</i> (95% CI)
	Mean	SD	Mean	SD						
Peak adductor longus force (BW)	0.74	0.38	0.54	0.27	2.57	0.19	0.04 - 0.34	0.012	0.17	0.04 - 0.74
Peak extensor hallucis longus force (BW)	0.22	0.09	0.28	0.09	2.77	0.06	0.02 - 0.10	0.007	724.81	7.72 - 111369.08
Flexor digitorum longus impulse (BW.ms)	10.03	5.46	7.03	2.88	2.77	2.70	0.76 - 4.64	0.007	0.83	0.71 - 0.968
Flexor hallucis longus impulse (BW.ms)	10.76	6.95	7.57	3.55	2.56	3.20	0.72 - 5.69	0.012	0.86	0.76 - 0.99
Gluteus minimus impulse (BW.ms)	130.34	35.28	115.47	24.33	2.75	18.22	5.02 - 31.42	0.007	0.98	0.96 - 1.00
Peak lateral gastrocnemius force (BW)	0.86	0.24	0.96	0.19	2.13	0.10	0.01 - 0.20	0.036	11.22	1.10 - 114.35
Peroneus tertius impulse (BW.ms)	8.22	3.30	5.84	2.62	3.07	1.96	0.69 - 3.24	0.003	0.77	0.64 - 0.94
Rectus femoris impulse (BW.ms)	166.05	76.86	225.19	52.79	3.87	57.54	27.91 - 87.16	<0.001	1.01	1.01 - 1.02
Peak semimembranosus force (BW)	0.52	0.13	0.65	0.20	3.06	0.11	0.04 - 0.18	0.003	90.44	3.75 - 2184.09
Semitendinosus impulse (BW.ms)	21.01	7.18	12.30	8.53	4.81	8.56	5.02 - 12.11	<0.001	0.88	0.82 - 0.94
Peak semitendinosus force (BW)	0.31	0.07	0.24	0.12	3.00	0.07	0.02 - 0.11	0.004	0.01	0.00 - 0.14

Soleus impulse (BW.ms)	467.68	118.87	396.30	80.47	3.33	76.05	30.62 - 121.48	0.001	0.99	0.99 - 1.00
Biceps femoris short head (%)	30.62	8.92	34.42	11.11	2.10	4.69	0.24 - 9.13	0.039	1.055	1.00 - 1.11
Semitendinosus (%)	15.68	4.60	9.51	4.62	5.70	5.89	3.83 - 7.94	<0.001	0.768	0.68 - 0.87
Rectus femoris (%)	19.49	4.97	25.97	7.25	4.64	6.52	3.72 - 9.31	<0.001	1.260	1.11 - 1.43
Vastus intermedius (%)	23.38	1.36	21.50	2.02	4.84	1.88	1.11 - 2.66	<0.001	0.405	0.25 - 0.65
Vastus lateralis (%)	35.95	2.78	33.20	3.53	3.88	2.79	1.36 - 4.21	<0.001	0.721	0.59 - 0.88
Vastus medialis (%)	21.17	1.35	19.33	1.83	5.07	1.85	1.12 - 2.57	<0.001	0.36	0.22 - 0.61
Mean co-contraction ratio	0.19	0.09	0.15	0.07	2.06	0.04	0.001 - 0.07	0.043	0.002	0.00 - 0.86

Table 5: Three-dimensional kinematics in male PFP participants and controls.

		Male control		Male PFP							
		Mean	SD	Mean	SD	<i>t</i>	<i>b</i>	<i>b</i> (95% CI)	<i>p</i>	<i>OR</i>	<i>OR</i> (95% CI)
Coronal plane	Pelvic ROM (°)	1.62	1.40	2.88	1.99	2.34	0.57	0.08 - 1.06	0.023	1.51	1.03 - 2.20
	Ankle angle at footstrike (°)	5.35	4.72	-4.06	5.21	6.85	4.83	3.41 - 6.25	p<0.001	0.67	0.54 - 0.83
	Peak ankle eversion (°)	-8.05	3.48	-14.69	6.13	5.07	3.53	2.13 - 4.93	p<0.001	0.68	0.54 - 0.86
Transverse plane	Ankle angle at footstrike (°)	-8.77	8.15	-15.33	5.49	3.40	3.35	1.37 - 5.33	0.001	0.86	0.78 - 0.96
	Ankle ROM (°)	7.24	3.91	0.69	1.12	8.02	3.24	2.43 - 4.04	p<0.001	0.37	0.22 - 0.64

Table 6: Muscle force indices in male PFP participants and controls.

		Male control		Male PFP							
		Mean	SD	Mean	SD	<i>t</i>	<i>b</i>	<i>b</i> (95% CI)	<i>p</i>	<i>OR</i>	<i>OR</i> (95% CI)
Adductor magnus impulse (BW.ms)		53.33	30.55	36.75	19.79	2.61	9.27	2.13 - 16.41	0.012	0.97	0.94 - 1.00
Gluteus minimus impulse (BW.ms)		126.55	38.37	108.45	24.42	2.32	10.28	1.37 - 19.19	0.025	0.98	0.96 - 1.01
Peroneus brevis impulse (BW.ms)		51.55	24.23	37.05	23.70	2.33	7.88	1.07 - 14.69	0.024	0.97	0.95 - 0.99
Peak peroneus brevis force (BW)		0.60	0.21	0.46	0.22	2.27	0.07	0.01 - 0.13	0.028	0.04	0.004 - 0.85

Semitendinosus impulse (BW.ms)	20.27	7.44	9.98	7.95	4.66	5.15	0.293 - 7.38	<0.001	0.85	0.77 - 0.93
Peak semitendinosus force (BW)	0.30	0.08	0.20	0.11	3.67	0.05	0.02 - 0.08	0.001	0.003	0.001 - 0.03
Soleus impulse (BW.ms)	464.90	135.71	383.28	73.12	2.74	42.16	11.22 - 73.10	0.009	0.99	0.98 - 1.00
Semitendinosus (%)	15.53	4.18	8.87	4.77	5.06	3.24	1.95 - 4.52	<0.001	0.74	0.63 - 0.88
Rectus femoris (%)	21.30	3.80	28.08	7.69	3.75	3.28	1.52 - 5.04	<0.001	1.321	1.10 - 1.59
Vastus intermedius (%)	23.04	1.13	20.98	2.18	4.01	1.01	0.50 - 1.51	<0.001	0.330	0.16 - 0.67
Vastus lateralis (%)	34.72	2.08	32.02	3.58	3.07	1.29	0.45 - 2.14	0.004	0.678	0.51 - 0.91
Vastus medialis (%)	20.94	1.07	18.92	2.04	4.18	0.99	0.51 - 1.46	<0.001	0.283	0.13 - 0.63

Table 7: Three-dimensional kinematics in female PFP participants and controls.

		Female control		Female PFP							
		Mean	SD	Mean	SD	<i>t</i>	<i>b</i>	<i>b</i> (95% CI)	<i>p</i>	<i>OR</i>	<i>OR</i> (95% CI)
Sagittal plane	Ankle angle at footstrike (°)	3.36	10.78	10.36	4.67	2.05	3.15	0.02 - 6.29	0.049	1.18	1.00 - 1.38
	Peak dorsiflexion (°)	16.51	4.29	20.19	3.12	2.73	1.88	0.05 - 3.29	0.01	1.35	1.04 - 1.74
Coronal plane	Pelvic ROM (°)	1.88	1.54	3.64	2.13	2.68	0.90	0.21 - 1.59	0.011	1.64	1.04 - 2.58
	Knee angle at footstrike (°)	1.16	5.38	-4.62	3.82	3.30	2.84	1.09 - 4.59	0.002	0.76	0.61 - 0.94
	Ankle angle at footstrike (°)	6.09	5.55	-4.93	7.41	4.39	5.10	2.73 - 7.47	P<0.001	0.75	0.60 - 0.93
	Peak ankle eversion (°)	-8.28	5.11	-15.75	9.32	2.49	3.30	0.60 - 6.00	0.018	0.86	0.75 - 0.99
Transverse plane	Knee angle at footstrike (°)	-12.32	7.83	-1.95	7.06	3.64	4.95	2.18 - 7.72	0.001	1.22	1.04 - 1.42
	Knee ROM (°)	17.88	5.82	10.39	3.88	3.76	3.34	1.58 - 5.15	0.001	0.75	0.60 - 0.94

Table 8: Joint loading and temporal indices in female PFP participants and controls.

	Female control		Female PFP		<i>t</i>	<i>b</i>	<i>b</i> (95% CI)	<i>p</i>	<i>OR</i>	<i>OR</i> (95% CI)
	Mean	SD	Mean	SD						
Patellofemoral impulse (BW.ms)	655.07	192.48	817.74	125.04	2.51	74.88	14.18 - 135.59	0.017	1.01	1.00 - 1.02
Peak patellofemoral force (BW)	6.13	1.79	7.31	1.08	2.13	0.58	0.02 - 1.14	0.041	1.73	1.01 - 2.98
Peak patellofemoral stress (KPa/BW)	10.82	2.86	13.07	1.71	2.54	1.11	0.22 - 2.00	0.016	1.53	1.06 - 2.21
Patellofemoral stress loading rate (KPa/BW/s)	318.53	124.27	402.55	129.66	2.52	55.48	10.60 - 100.36	0.017	1.01	1.00 - 1.02
Patellofemoral stress impulse (KPa/BW.ms)	1205.92	314.27	1567.44	192.80	3.50	167.22	70.00 - 264.45	0.001	1.00	1.00 - 1.02
Patellofemoral stress per mile (BW)	764.95	429.71	1116.67	361.68	2.64	203.77	50.07 - 357.46	0.01	1.01	1.00 - 1.02
Vertical GRF impulse (BW.ms)	309.02	34.06	354.18	37.04	3.17	20.13	7.21 - 33.06	0.003	1.04	1.01 - 1.07
Strike index (%)	40.20	21.69	15.49	3.37	3.94	11.29	5.45 - 17.13	p<0.001	0.79	0.65 - 0.95

Table 9: Muscle force indices in female PFP participants and controls.

	Female control		Female PFP		<i>t</i>	<i>b</i>	<i>b</i> (95% CI)	<i>p</i>	<i>OR</i>	<i>OR</i> (95% CI)
	Mean	SD	Mean	SD						
Peak adductor longus force (BW)	0.75	0.41	0.51	0.24	2.07	0.13	0.06 - 0.16	0.043	0.21	0.01 - 0.97
Peak biceps femoris force (BW)	0.80	0.23	1.01	0.28	2.52	0.12	0.02 - 0.21	0.017	55.52	1.47 - 2091.52
Peak extensor hallucis longus force (BW)	0.17	0.06	0.27	0.12	2.89	0.05	0.01 - 0.09	0.007	215238	4.57 - 10134951829.16
Peak lateral gastrocnemius force (BW)	0.81	0.24	1.02	0.18	2.59	0.10	0.02 - 0.18	0.014	199.27	2.16 - 18373.01
Peak medial gastrocnemius force (BW)	1.79	0.24	2.12	0.33	3.38	0.17	0.07 - 0.28	0.002	305.1	2.72 - 34221.81
Peak peroneus longus force (BW)	0.95	0.25	1.29	0.37	3.34	0.19	0.07 - 0.30	0.002	49.52	2.37 - 1035.09
Psoas major impulse (BW.ms)	83.49	35.93	125.46	45.12	3.07	22.54	7.60 - 37.48	0.004	1.03	1.00 - 1.06
Rectus femoris impulse (BW.ms)	117.08	49.13	199.54	49.23	4.69	41.97	23.72 - 60.21	p<0.001	1.04	1.01 - 1.07
Peak semimembranosus force (BW)	0.52	0.15	0.75	0.19	3.36	0.09	0.04 - 0.15	0.002	2323.7	5.15 - 1047595.46
Total quadriceps force impulse (BW.ms)	675.62	171.35	891.41	117.88	3.78	101.72	46.84 - 156.60	0.001	1.01	1.00 - 1.02
Peak total quadriceps force (BW)	5.88	1.52	7.18	1.05	2.68	0.64	0.15 - 1.14	0.012	2.05	1.07 - 3.92
Vastus intermedius impulse (BW.ms)	160.63	40.36	199.77	29.64	2.75	17.73	4.59 - 30.86	0.01	1.03	1.00 - 1.05

Peak vastus intermedius impulse (BW.ms)	1.64	0.48	2.02	0.32	2.52	0.19	0.04 - 0.34	0.017	9.05	1.16 - 70.48
Vastus medialis impulse (BW.ms)	145.29	38.61	178.72	27.68	2.48	15.24	2.73 - 27.75	0.019	1.03	1.00 - 1.06
Peak vastus medialis force (BW)	1.50	0.45	1.83	0.31	2.32	0.17	0.02 - 0.31	0.027	8.48	1.02 - 70.87
Semitendinosus (%)	15.88	5.23	10.59	4.30	2.67	2.29	0.54 - 4.05	0.012	0.80	0.63 - 0.98
Rectus femoris (%)	17.06	5.42	22.46	4.88	3.21	2.99	1.09 - 4.89	0.003	1.27	1.05 - 1.54
Vastus intermedius (%)	23.84	1.53	22.38	1.39	2.99	0.79	0.25 - 1.33	0.005	0.45	0.23 - 0.88
Vastus lateralis (%)	37.62	2.79	35.17	2.46	3.00	1.41	0.45 - 2.38	0.005	0.66	0.47 - 0.93
Vastus medialis (%)	21.49	1.64	20.00	1.20	2.97	0.79	0.25 - 1.32	0.006	0.45	0.23 - 9.90

		Male PFP		Female PFP		<i>t</i>	<i>b</i>	<i>b</i> (95% CI)	<i>p</i>
		Mean	SD	Mean	SD				
Sagittal plane	Hip angle at footstrike (°)	31.30	8.46	39.54	7.02	3.55	8.79	3.79 - 13.79	0.001
	Knee angle at footstrike (°)	10.15	4.69	14.69	5.49	2.89	4.62	1.38 - 7.86	0.006
	Peak knee flexion (°)	38.32	8.51	43.59	5.17	2.54	5.83	1.19 - 10.48	0.015
Coronal plane	Peak hip adduction (°)	7.62	5.73	11.30	2.71	2.38	3.71	0.57 - 6.85	0.02
Transverse plane	Knee ROM (°)	14.53	4.80	10.39	3.88	2.75	3.80	1.00 - 6.59	0.009

Table 10: Three-dimensional kinematics in male and female PFP participants.

Table 11: Joint loading and temporal indices in male and female PFP participants.

	Male PFP		Female PFP		<i>t</i>	<i>b</i>	<i>b</i> (95% CI)	<i>p</i>
	Mean	SD	Mean	SD				
Loading rate (BW/s)	150.08	45.89	173.28	39.12	2.18	27.91	1.97 - 53.85	0.036

Table 12: Muscle force indices in male and female PFP participants.

	Male PFP		Female PFP		<i>t</i>	<i>b</i>	<i>b</i> (95% CI)	<i>p</i>
	Mean	SD	Mean	SD				
Biceps femoris long head (BW.ms)	21.13	11.36	31.88	16.29	2.55	10.96	2.28 - 19.64	0.015
Peak biceps femoris short head force (BW)	0.81	0.31	1.01	0.28	2.45	0.22	0.04 - 0.41	0.019
Flexor digitorum longus impulse (BW.ms)	47.30	24.94	28.09	12.48	2.69	17.73	4.41 - 31.04	0.01
Flexor hallucis longus impulse (BW.ms)	6.40	2.78	8.06	2.82	2.29	1.93	0.23 - 3.63	0.027
Peak gluteus maximus force (BW)	142.04	48.48	178.45	45.06	2.47	37.41	6.82 - 67.99	0.018
Gluteus maximus impulse (BW.ms)	1.40	0.45	1.69	0.42	2.81	0.35	0.10 - 0.60	0.008
Gluteus medius impulse (BW.ms)	261.79	58.62	320.89	48.67	3.24	54.01	20.31 - 87.72	0.002
Peak gluteus medius force (BW)	2.25	0.45	2.66	0.46	2.76	0.39	0.11 - 0.68	0.009
Gluteus minimus impulse (BW.ms)	108.45	24.42	127.16	19.81	2.43	17.46	2.94 - 31.98	0.02
Peak gluteus minimus force (BW)	0.95	0.21	1.16	0.22	3.00	0.20	0.07 - 0.34	0.005
Peak gracilis force (BW)	0.08	0.05	0.11	0.04	2.93	0.04	0.01 - 0.06	0.006
Peak iliacus force (BW)	2.01	0.39	2.40	0.37	3.48	0.42	0.17 - 0.66	0.0012
Rectus femoris impulse (BW.ms)	240.58	49.58	199.54	49.23	2.55	40.50	8.42 - 72.58	0.015
Peak sartorius force (BW)	0.24	0.05	0.30	0.05	3.76	0.06	0.03 - 0.09	0.001
Peak semimembranosus impulse (BW.ms)	36.65	15.54	55.55	19.95	3.30	17.59	6.80 - 28.38	0.002
Peak semimembranosus force (BW)	0.58	0.17	0.75	0.19	2.91	0.16	0.05 - 0.27	0.0059
Semitendinosus impulse (BW.ms)	9.98	7.95	16.17	8.28	2.54	6.51	1.32 - 11.70	0.015
Peak semitendinosus force (BW)	0.20	0.11	0.30	0.13	2.96	0.11	0.03 - 0.18	0.005
Tensor fasciae latae impulse (BW.ms)	55.82	10.98	66.03	9.17	3.04	10.08	3.37 - 16.80	0.004
Peak tensor fasciae latae force (BW)	0.45	0.08	0.57	0.07	4.98	0.12	0.07 - 0.16	p<0.001
Tibialis impulse (BW.ms)	25.39	14.36	34.54	11.76	2.94	10.99	3.44 - 18.55	0.005
Total hamstring impulse (BW.ms)	108.48	36.77	147.66	45.69	3.02	39.05	12.90 - 65.21	0.004
Peak total hamstring force (BW)	1.39	0.47	1.66	0.46	2.01	0.29	0.06 - 0.59	0.048
Rectus femoris (%)	28.08	7.69	22.46	4.88	2.67	5.79	1.40 - 10.18	0.011

Vastus intermedius (%)	20.98	2.18	22.38	1.39	2.38	1.46	0.22 - 2.70	0.022
Vastus lateralis (%)	32.02	3.58	35.17	2.46	3.10	3.20	1.11 - 5.28	0.004
Mean co-contraction ratio	0.13	0.07	0.19	0.06	2.34	0.05	0.006 - 0.09	0.024
Peak co-contraction ratio	0.19	0.08	0.25	0.09	2.12	0.06	0.003 - 0.12	0.04